

# Street Tree Assessment Report

## *Winchester, Virginia*

### Overview

Street trees are a vital community asset that enhance our day-to-day lives and mitigate many of the negative impacts of urbanization. In 2008, a sample street tree inventory was conducted in Winchester, Virginia to assess tree abundance, composition, functional benefits, and monetary value. Trees residing within the right-of-way along 10% of public streets were surveyed to determine their species, size, condition, and placement. Inventory data were collected by Virginia Tech for this assessment report. The inventory data were analyzed using i-Tree Streets assessment software developed by the U.S. Forest Service.

### Key Findings

- Winchester has an estimated 8,990 street trees.
- Winchester's five most abundant street tree species are sugar maple, Norway maple, American sycamore, Callery pear, and red maple.
- Each year, Winchester's street trees intercept over 26 million gallons of rainfall and sequester over 3 million pounds of carbon dioxide.
- In total, Winchester's street trees provide about \$980 thousand in benefits annually or roughly \$109 per tree.
- The replacement value of Winchester's street trees is estimated at over \$33 million.

*Prepared by Virginia Tech  
Department of Forest Resources and  
Environmental Conservation*

<http://urbanforestry.frec.vt.edu/>

[arborist@vt.edu](mailto:arborist@vt.edu)



 **Virginia Tech**  
*Invent the Future®*



## Street Tree Abundance and Composition

Winchester's estimated street tree population is 8,990. Winchester's street trees provide about 159 acres of canopy, which covers roughly 2.7% of the land area. The five most abundant species are sugar maple (13%), Norway maple (12%), American sycamore (8%), Callery pear (7%), and red maple (7%). The most important species (accounting for leaf area and canopy cover in addition to tree count) include sugar maple (21%), Norway maple (13%), American sycamore (12%), Siberian elm (6%), and Callery pear (5%).

Large-stature, broadleaf deciduous trees are the most common tree form amongst Winchester's street trees. About 50% of Winchester's street trees are smaller than 12 in. trunk diameter while less than 5% are larger than 30 in. The majority of Winchester's street trees (~96%) were rated in fair to good condition.



### Relative abundance of Winchester's street trees by foliage type and mature height class.

Foliage Type	Small (< 25')	Medium (25 - 45')	Large (> 45')	Total	% of Total
Broadleaf Deciduous	1,919	1,283	5,082	8,284	92
Broadleaf Evergreen	99	20	0	119	1
Conifer Evergreen	20	448	119	587	7
<b>Total</b>	<b>2,038</b>	<b>1,751</b>	<b>5,201</b>	<b>8,990</b>	<b>100</b>
<b>% of Total</b>	<b>23</b>	<b>19</b>	<b>58</b>	<b>100</b>	

## Street Tree Benefits and Value

Gross annual benefits provided by Winchester’s street trees are valued at \$983,585. These benefits come from contributions that street trees make to real estate aesthetics, rainfall interception, energy conservation, air pollution reduction, and CO<sub>2</sub> sequestration. Each year, Winchester’s street trees intercept roughly 26 million gallons of rainfall, conserve a combined 803 megawatt-hour of electricity and 29 thousand therms of natural gas for home cooling and heating, and remove about 3.4 million pounds of carbon from the atmosphere. In addition, Winchester’s street trees currently store about 40 million pounds of carbon, which is valued at over \$300 thousand. Although Winchester’s street trees have a net positive impact on air pollution – removing over 1,703 pounds of pollutants annually – its current mix of tree species heavily emits biogenic volatile organic compounds (BVOCs), which results in a negative monetary value for pollution reduction.

On a per-tree basis, the most beneficial tree species are Siberian elm (\$234 per year), silver maple (\$189

per year), sugar maple (\$187 per year), sweetgum (\$178 per year), and American sycamore (\$170 per year). These values reflect the large size that these trees have attained, providing abundant leaf area and canopy cover. The average street tree provides \$109 in gross benefits annually. Gross benefits do not account for annual costs associated with planting, maintenance, or removal, which were not available for this analysis.

The replacement value of Winchester’s street tree population is estimated at \$33,228,972. This is the value of street trees as a structural asset, and reflects the cost to replant trees in a quantity sufficient to replace their current level of functional benefits. Because a large street tree produces the same amount of benefits as numerous nursery-sized trees, replacing a large tree would require significant resources that may not be feasible due to both spatial and budgetary constraints.

### Gross annual benefits provided by Winchester's street trees.

Benefit Type	Resource Units	Total \$	Avg. \$/Tree
Aesthetic enhancements	–	613,838	68.28
Rainfall Interception (gallons)	26,445,189	261,826	29.12
Energy Conservation <sup>1</sup>	–	91,015	10.12
Electricity (MWh)	803	60,962	–
Natural Gas (therms)	28,732	30,530	–
Air Pollution reduction (lb) <sup>2</sup>	1,703	-11,525	-1.28
CO <sub>2</sub> sequestration (lb) <sup>3</sup>	3,410,385	25,578	2.85
<b>Total Benefits</b>	–	<b>980,732</b>	<b>109.09</b>

<sup>1</sup>Sum of electricity and natural gas conservation.

<sup>2</sup>Net pollution reduction (O<sub>3</sub>, NO<sub>2</sub>, PM<sub>10</sub>, and SO<sub>2</sub>) accounting for pollutant deposition, pollutant avoidance, and BVOC emissions. Note, if Resource Units value is negative, BVOC emissions exceeded pollution reduction. If only total \$ is negative, then BVOC pricing exceeded pollutant pricing, but pollution reduction still occurred.

<sup>3</sup>Net sequestration accounting for gross tree sequestration, tree decomposition emissions, and tree maintenance machinery emissions.

## Street Tree Opportunities

Winchester has a highly valuable street tree population. To sustain this resource and its benefits, the city should continue to focus on planting diverse, functional species and maintaining trees to ensure their health, safety, and appearance. Urban forestry experts generally recommend that a municipal tree population comprise no more than 10% of a single species and 20% of a single genus in order to minimize impacts of pest outbreaks and other species-specific disorders. Sugar maple (13%) and Norway maple (12%) both exceed the species threshold. Collectively, maple species account for 34% of the street trees. Although maples are proven performers, planting efforts should temper their use to ensure the diversity and health of Winchester's street trees.

One of the most noxious pests threatening Virginia's street trees is emerald ash borer, an insect introduced from Asia that has killed millions of native ash trees in the United States. Fortunately, native ash species comprise just 5.8% of Winchester's street trees and account for only 4.7% of the street tree canopy cover. However, Winchester must remain vigilant in managing street tree diversity because there is ongoing risk of unforeseen introduction of noxious tree pests into the United States.

About 77% of Winchester's street tree population comprises medium- and large-stature species such as maple and oak. This is a favorable distribution given that larger trees provide higher levels of benefits, yet presence of overhead utility lines may require planting of small-stature tree species in certain places to minimize power disruptions and pruning costs.

The size distribution of Winchester's street trees suggests a stable age structure. Because street trees inevitably grow old and die or must be removed to accommodate

land use changes, an ample number of young trees must always exist in order to sustain street tree benefits. The fact that the two diameter classes that encompass the largest percentage of the total street tree population are the 0-6 and 6-12 inch diameter classes, respectively, is a source of optimism. However, ongoing planting efforts, with particular focus on large stature, highly functional tree species, should be taken to ensure a high level of benefits will be provided by Winchester's street trees for the future.

Winchester's street trees comprise a number of species that produce large amounts of BVOCs, which are precursors to ground-based ozone. Heavy emitters of BVOCs in Winchester include Siberian elm, sweetgum, and American sycamore. Winchester should consider planting more low-BVOC street trees such as ginkgo, linden, and certain maples if maximizing air quality benefits is a key community objective. However, this planting strategy should not compromise efforts to maximize canopy cover or species diversity. Urban forestry experts generally believe that trees have a net positive impact on air quality, regardless of BVOC emissions, by lowering air temperature and reducing fossil fuel combustion in urban areas.

This assessment has reported gross benefits of Winchester's street trees, which may not fully reflect the true value of this vital resource. Direct and indirect costs of administering and managing street trees can vary considerably based on species composition, tree size distribution, and other local environmental and economic factors. Therefore, findings of this report should be carefully interpreted in the context of local circumstances that impact tree benefits and costs.

## About This Report

This report was co-authored by Eric Wiseman and Julia Bartens with the [Department of Forest Resources and Environmental Conservation](#) at Virginia Tech. Report layout and design by Sarah Gugercin.

This report was made possible through grants from the Virginia Department of Forestry and the U.S. Forest Service. Technical assistance was graciously provided by the Davey Resource Group.

Inventory data were analyzed using i-Tree Streets assessment software version 4.0.4. Benefit estimates were based on i-Tree modeling data from the Charlotte, North Carolina reference city in the South Climate Zone. The 2010 median home price, used to calculate street tree aesthetic benefits for Winchester was \$267,000 as reported by the U.S. Census Bureau in <http://quickfacts.census.gov/qfd/index.html>. Additional information about methods used in this street tree assessment can be found [on our website](#).

Date of Publication: July 2012.